



SUMMARY OF ANTIMICROBIAL ACTIVITY

ANIMAL FACILITY CONCENTRATED DISINFECTANT CLEANER & DEODORIZER

Cleaner – Disinfectant – Virucide*

Description

Animal Facility Concentrated Disinfectant is a broad spectrum, moderate pH, hard surface disinfectant. When used as directed, this product will deliver effective biocidal action against bacteria, fungi, and viruses. This formulation is a blend of a premium active ingredients and inerts: surfactants, chelates, and water. Biocidal performance is attained when this product is properly diluted at 1 oz. per gallon or 1:128. (See below for additional dilutions and corresponding efficacy).

Animal Facility Concentrated Disinfectant can be in Veterinary clinics, animal life science laboratories, animal research centers, animal quarantine areas, animal breeding facilities, animal holding areas, kennels, dog/cat animal kennels, breeding and grooming establishments, pet animal quarters, zoos, pet shops, tack shops, operating rooms, washing areas, waiting rooms, examination rooms, and other animal care facilities.

This product can be used to disinfect, clean and deodorize terrarium and small animal cages, hot rock, substrate and cage furniture, plastic terrarium ornaments, driftwood, heat caves and water dishes.

Regulatory Summary

Physical Properties

EPA Registration No. 10324-105-86550 USDA Authorization None California Status Canadian PCP# None Canadian Din # None	pH of Concentrate 9.5 – 11.5 Specific Gravity @ 25°C 1.06 Pounds per gallon @ 25°C 8.79	Flash Point (PMCC) >200°F % Quat (mol. wt.360.5) 9.0 – 9.45 % Volatile 85+
--	--	---

Efficacy For ProVetLogic Professional Animal Facility Disinfectant

Hospital and General Disinfection

This product is bactericidal according to the AOAC Use Dilution Test method on hard inanimate surfaces modified in the presence of 5% organic serum at 1 ounce of this product per gallon of water (703 ppm active). Treated surfaces must remain wet for 10 minutes.

(Testing is performed per the AOAC UDT/GST method (DIS/TSS-1). Sixty carriers are required on 3 separate lots, one of which must be > 60 days old against *Pseudomonas aeruginosa*, *Salmonella enterica* and *Staphylococcus aureus*. Killing of 59 out of 60 carriers is required (total carriers = 540).)

CARRIER	POPULATION	SAMPLE #	CARRIERS	# POSITIVE
<i>Pseudomonas aeruginosa</i> ATCC #15442	8.2 X 10 ⁶ CFU/Carrier	A (60 Days Old)	60	0/60
	4.3 X 10 ⁶ CFU/Carrier	B	60	1/60
	9.7X 10 ⁵ CFU/Carrier	C	60	1/60
<i>Salmonella enterica</i> ATCC #10708	1.4 X 10 ⁴ CFU/Carrier	A (60 Days Old)	60	0/60
	7.3 X 10 ⁴ CFU/Carrier	B	60	0/60
	4.8 X 10 ⁵ CFU/Carrier	C	60	0/60
<i>Staphylococcus aureus</i> ATCC #6538	1.4 X 10 ⁶ CFU/Carrier	A (60 Days Old)	60	1/60
	2.8 X 10 ⁵ CFU/Carrier	B	60	1/60
	9.2 X 10 ⁵ CFU/Carrier	C	60	1/60

Supplemental Organisms

(Testing is performed per the AOAC UDT/GST method. Ten carriers are required on 2 separate lots against each supplemental organism. Killing of 10 out of 10 carriers is required (total carriers = 20).)

CARRIER	POPULATION	SAMPLE #	CARRIERS	# POSITIVE
<i>Campylobacter jejuni</i> ATCC 29428	4.0 X 10 ⁴ CFU/Carrier	A	10	0/10
		B	10	0/10
<i>Corynebacterium ammoniagenes</i> ATCC 6871	9.6 X 10 ⁴ CFU/Carrier	A	10	0/10
		B	10	0/10
<i>Enterobacter aerogenes</i> ATCC 13048	4.1 X 10 ⁵ CFU/Carrier	A	10	0/10
		B	10	0/10
<i>Enterococcus faecalis</i> ATCC 19433	3.5 X 10 ⁶ CFU/Carrier	A	10	0/10
		B	10	0/10
<i>Escherichia coli</i> ATCC 11229	3.5 X 10 ⁵ CFU/Carrier	A	10	0/10
		B	10	0/10
<i>Klebsiella pneumoniae</i> ATCC 13883	4.2 X 10 ⁴ CFU/Carrier	A	10	0/10
		B	10	0/10
<i>Listeria monocytogenes</i> ATCC 984	4.5 X 10 ⁵ CFU/Carrier	A	10	0/10
		B	10	0/10
<i>Salmonella schottmuelleri</i> ATCC 8759	9.6 X 10 ⁶ CFU/Carrier	A	10	0/10
		B	10	0/10
<i>Salmonella typhi</i> ATCC 6539	1.7 X 10 ⁵ CFU/Carrier	A	10	0/10
		B	10	0/10
<i>Shigella dysenteriae</i> ATCC 9361	3.7 X 10 ⁵ CFU/Carrier	A	10	0/10
		B	10	0/10
<i>Staphylococcus aureus</i> (Methicillin Resistant) (MRSA) ATCC 33591	1.3 X 10 ⁵ CFU/Carrier	A	10	0/10
		B	10	0/10
<i>Staphylococcus aureus</i> Community Associated Methicillin Resistant (CA-MRSA)	1.76 X 10 ⁵ CFU/Carrier	A	10	0/10
		B	10	0/10
<i>Streptococcus salivarius</i> ATCC 13419	1.5 X 10 ⁴ CFU/Carrier	A	10	0/10
		B	10	0/10

Virucidal against (at 1 ounce)

This product was evaluated at 1 ounce per gallon use level (703 ppm quat active), in the presence of 5% serum with a 10 minute contact time and found to be effective against the following viruses on hard nonporous environmental surfaces.

(Testing is performed per EPA Guidance (DIS/TSS-7). Two separate lots are tested. Inactivation of virus must be demonstrated at all dilutions when no cytotoxicity is observed or at all dilutions above the cytotoxic level when it is observed. The data must demonstrate a 3-log reduction in viral titer for both lots.)

	DRIED VIRUS CONTROL	SAMPLE	RESULT	LOG REDUCTION
Avian Influenza A (H5N1) virus	4.5 Log ₁₀	A	0.5 Log ₁₀	4.0 Log ₁₀
		B	0.5 Log ₁₀	4.0 Log ₁₀
Avian influenza/Turkey/Wisconsin ATCC VR-798	7.5 Log ₁₀	A	1.8 Log ₁₀	5.7 Log ₁₀
		B	1.8 Log ₁₀	5.7 Log ₁₀
Avian Reovirus ATCC VR-2449	6.0 Log ₁₀	A	0.5 Log ₁₀	5.5 Log ₁₀
		B	0.5 Log ₁₀	5.5 Log ₁₀
Bovine Viral Diarrhea ATCC VR-534	4.5 Log ₁₀	A	0.5 Log ₁₀	4.0 Log ₁₀
		B	0.5 Log ₁₀	4.0 Log ₁₀
Canine Distemper ATCC VR-64	4.8 Log ₁₀	A	1.5 Log ₁₀	3.3 Log ₁₀
		B	1.5 Log ₁₀	3.3 Log ₁₀
Canine Coronavirus ATCC VR-809	4.5 Log ₁₀	A	1.5 Log ₁₀	3.0 Log ₁₀
		B	0.5 Log ₁₀	4.0 Log ₁₀
Equine Arteritis virus ATCC VR-796	5.75 Log ₁₀	A	0.5 Log ₁₀	5.25 Log ₁₀
		B	0.5 Log ₁₀	5.25 Log ₁₀
Hepatitis B Virus	6.84 Log ₁₀	A	1.78 Log ₁₀	5.06 Log ₁₀
		B	2.05 Log ₁₀	4.79 Log ₁₀
		Confirmatory B	2.35 Log ₁₀	4.79 Log ₁₀
Hepatitis C Virus ATCC CCL-22	6.85 Log ₁₀	A	1.29 Log ₁₀	5.56 Log ₁₀
		B	1.06 Log ₁₀	5.79 Log ₁₀
		Confirmatory B	1.06 Log ₁₀	.79 Log ₁₀
Herpes Simplex Type1 ATCC VR-266	6.8 Log ₁₀	A	1.8 Log ₁₀	5.0 Log ₁₀
		B	1.8 Log ₁₀	5.0 Log ₁₀
Human Coronavirus ATCC VR-740	4.5 Log ₁₀	A	1.5 Log ₁₀	3.0 Log ₁₀
		B	1.5 Log ₁₀	3.0 Log ₁₀
Human Immunodeficiency Virus type 1 (HIV 1) HTLV-IIIB	6.0 Log ₁₀	A	2.5 Log ₁₀	3.5 Log ₁₀
		B	2.5 Log ₁₀	3.5 Log ₁₀
Infectious Bovine Rhinotracheitis virus (IBR) ATCC VR-188	5.2 Log ₁₀	A	1.5 Log ₁₀	3.7 Log ₁₀
		B	1.5 Log ₁₀	3.7 Log ₁₀
Infectious Bronchitis Virus Beaudette IB42	5.25 Log ₁₀	A	0.5 Log ₁₀	4.75 Log ₁₀
		B	0.5 Log ₁₀	4.75 Log ₁₀
Infectious Laryngotracheitis Virus (LT) Strain LT-IVAX	4.5 Log ₁₀	A	0.5 Log ₁₀	4.0 Log ₁₀
		B	0.5 Log ₁₀	4.0 Log ₁₀
Influenza A2/Japan/305 ATCC VR-100	7.5 Log ₁₀	A	1.8 Log ₁₀	5.7 Log ₁₀

	DRIED VIRUS CONTROL	SAMPLE	RESULT	LOG REDUCTION
Newcastle disease virus	6.0 Log ₁₀	B	1.8 Log ₁₀	5.7 Log ₁₀
		A	1.8 Log ₁₀	4.2 Log ₁₀
Norwalk/Norovirus ATCC VR-782	5.75 Log ₁₀	B	1.8 Log ₁₀	.2 Log ₁₀
		A	1.5 Log ₁₀	4.25 Log ₁₀
Porcine Respiratory & Reproductive (PRRSV)	5.5 Log ₁₀	B	1.5 Log ₁₀	4.25 Log ₁₀
		Confirmatory A	0.5 Log ₁₀	5.0 Log ₁₀
Porcine Rotavirus ATCC VR-893	4.5 Log ₁₀	A	1.5 Log ₁₀	4.0 Log ₁₀
		B	1.5 Log ₁₀	4.0 Log ₁₀
Pseudorabies virus ATCC VR-135	4.5 Log ₁₀	A	1.5 Log ₁₀	3.0 Log ₁₀
		B	1.5 Log ₁₀	3.0 Log ₁₀
Transmissible Gastroenteritis (TGE) ATCC VR-742	5.7 Log ₁₀	A	1.5 Log ₁₀	3.0 Log ₁₀
		B	1.5 Log ₁₀	3.0 Log ₁₀
Vaccinia virus Hoffmann La Roche	6.8 Log ₁₀	A	2.5 Log ₁₀	3.2 Log ₁₀
		B	2.5 Log ₁₀	3.2 Log ₁₀
		A	2.5 Log ₁₀	4.3 Log ₁₀
		B	2.5 Log ₁₀	4.3 Log ₁₀

Virucidal against (at 1 1/4 ounce)

This product was evaluated at 1.25 ounces per gallon use level (879 ppm quat active), in the presence of 5% serum with a 10 minute contact time and found to be effective against the following viruses on hard nonporous environmental surfaces.

	DRIED VIRUS CONTROL	SAMPLE	RESULT	LOG REDUCTION
Canine Adenovirus ATCC VR-800	6.0 Log ₁₀	A	0.5 Log ₁₀	5.5 Log ₁₀
		B	1.5 Log ₁₀	4.5 Log ₁₀
Canine Parainfluenza Virus ATCC VR-666	4.75 Log ₁₀	A	0.5 Log ₁₀	4.25 Log ₁₀
		B	0.5 Log ₁₀	4.25 Log ₁₀
Feline Calicivirus ATCC VR-782	6.5 Log ₁₀	A	1.5 Log ₁₀	5.0 Log ₁₀
		B	1.5 Log ₁₀	5.0 Log ₁₀
Feline leukemia	6.5 Log ₁₀	A	2.5 Log ₁₀	4.0 Log ₁₀
		B	2.5 Log ₁₀	4.0 Log ₁₀
Feline Infectious Peritonitis ATCC VR-2202	5.25 Log ₁₀	A	1.5 Log ₁₀	3.75 Log ₁₀
		B	1.5 Log ₁₀	3.75 Log ₁₀
Feline Panleukopenia ATCC VR-648	4.5 Log ₁₀	A	0.5 Log ₁₀	4.0 Log ₁₀
		B	0.5 Log ₁₀	4.0 Log ₁₀
Feline Picornavirus ATCC VR-649	4.5 Log ₁₀	A	1.5 Log ₁₀	3.0 Log ₁₀
		B	1.5 Log ₁₀	3.0 Log ₁₀
Feline Rhinotrachetis ATCC VR-636	5.0 Log ₁₀	A	0.5 Log ₁₀	4.5 Log ₁₀
		B	0.5 Log ₁₀	4.5 Log ₁₀
Infectious Canine Hepatitis Virus ATCC VR-293	6.25 Log ₁₀	A	1.5 Log ₁₀	4.75 Log ₁₀
		B	1.5 Log ₁₀	4.75 Log ₁₀
Murine Parainfluenza virus type 1 ATCC VR-105	7.5 Log ₁₀	A	1.5 Log ₁₀	6.0 Log ₁₀
		B	1.5 Log ₁₀	6.0 Log ₁₀

	DRIED VIRUS CONTROL	SAMPLE	RESULT	LOG REDUCTION
Rabies Virus	6.23 Log ₁₀	A	2.5 Log ₁₀	3.73 Log ₁₀
		B	2.5 Log ₁₀	3.73 Log ₁₀

Virucidal against (at 4 ounces)

This product was evaluated at 4 ounces per gallon use level (2800 ppm quat active), in the presence of 5% serum with a 10 minute contact time and found to be effective against the following viruses on hard nonporous environmental surfaces.

	DRIED VIRUS CONTROL	SAMPLE	RESULT LOG	REDUCTION
Canine Parvovirus	7.5 Log ₁₀	A	4.0 Log ₁₀	3.5 Log ₁₀
		B	4.33 Log ₁₀	3.17 Log ₁₀
Minute virus of Mice (Parvovirus) ATCC VR-1346	5.75 Log ₁₀	A	2.5 Log ₁₀	3.25 Log ₁₀
		B	2.5 Log ₁₀	3.25 Log ₁₀
Porcine Parvovirus ATCC VR-742	5.5 Log ₁₀	A	2.5 Log ₁₀	3.0 Log ₁₀
		B	2.5 Log ₁₀	3.0 Log ₁₀

Fungicidal against

This product was evaluated at 1 ounce per gallon with a 10 minute contact time and found to be effective against the following fungi on hard nonporous environmental surfaces.

(Testing is performed per the AOAC fungicidal method (DIS/TSS-6). Two separate lots are tested against Trichophyton mentagrophytes in a suspension test. Killing of all fungal spores in 10 minutes is required.)

	CARRIER POPULATION	SAMPLE #	CARRIERS #	POSITIVE
<i>Trichophyton mentagrophytes</i> ATCC #9533	1.1 X 10 ⁶ CFU/Carrier	A	10	0/10
		B	10	0/10

Mold and Mildew Control

Use this product at 1 ounce per gallon to control the growth of mold and mildew and their odors on hard, non-porous surfaces. Thoroughly wet all treated surfaces completely. Let air-dry. Repeat application weekly or when growth or odor reappears.

	TILE NUMBER	UNTREATED AFTER 7 DAYS	SAMPLE A AFTER 7 DAYS	SAMPLE B AFTER 7 DAYS
<i>Aspergillus niger</i> ATCC #6275	1	Growth 80%	No Growth 0%	No Growth 0%
	2	Growth 100%	No Growth 0%	No Growth 0%
	3	Growth 80%	No Growth 0%	No Growth 0%
	4	Growth 80%	No Growth 0%	No Growth 0%
	5	Growth 80%	No Growth 0%	No Growth 0%
	6	Growth 80%	No Growth 0%	No Growth 0%
	7	Growth 80%	No Growth 0%	No Growth 0%
	8	Growth 100%	No Growth 0%	No Growth 0%
	9	Growth 100%	No Growth 0%	No Growth 0%
	10	Growth 80%	No Growth 0%	No Growth 0%

Non-Food Contact Surface Sanitizer

Add 1 ounce of this product to 1 gallon of water to sanitize hard porous and non-porous non-food contact surfaces. Treated surfaces must remain wet for 60 seconds. Then wipe with sponge, mop or cloth or allow to air dry. At this dilution food contact surfaces must be rinsed.

(Testing is performed per EPA Guidance (DIS/TSS-10). Three lots are required, one of which must be 60 days old. Testing is performed against *Staphylococcus aureus* and *Klebsiella pneumoniae* containing 5% organic load. *Enterobacter aerogenes* may be substituted for *Klebsiella pneumoniae*. The results must show a reduction of at least 99.9% in the number of each test microorganism over the parallel control count within 5 minutes.)

	CARRIER POPULATION	SAMPLE	SURVIVOR AVERAGE LOG₁₀	60 SECOND PERCENT KILL
<i>Klebsiella pneumoniae</i> ATCC 4352 (60 Days Old)	5.71 Log ₁₀	A	>4.23 Log ₁₀	>99.9
		B	>4.31 Log ₁₀	>99.9
		C	>4.03 Log ₁₀	>99.9
<i>Staphylococcus aureus</i> ATCC #6538 (60 Days Old)	6.45 Log ₁₀	A	>4.69 Log ₁₀	>99.9
		B	>4.87 Log ₁₀	>99.9
		C	>4.33 Log ₁₀	>99.9